



Travel Vaccination Schedule

Vaccine	Product Name	Doses	Dosing interval	Fee	Effective duration
Hepatitis A	Aimmugen	3 doses	0, 2 to 4 weeks later, 24 weeks later	JPY 9,200	more than 5 years
Hepatitis B	Heptavax, Bimmugen	3 doses	0, 4 weeks later, 20 to 24 weeks after the first shot	JPY 5,000	more than 10 years
MR	Freeze-dried live attenuated Measles and Rubella combined vaccine	2 doses	0, 4 weeks later	JPY 9,900	20 to 30 years
Rabies	*VERORAB	2 doses for pre-exposure	0, 7 days later	JPY 16,000	Life-long
		3 to 4 doses for post-exposure	【previously vaccinated】 0, 3 days later (total 2 doses) 【not previously vaccinated】 ①WHO (Essen) method: 0, 3, 7, 14 to 28 days later (total 4 doses) ②Zagreb method: 0, 7, 21 days later (total 3 doses) ③Thailand Red Cross intracutaneous method: 0, 3, 7 days later (total 3 doses)		
Typhoid	*Typhim Vi	1 dose	only once	JPY 10,000	3 years
Tetanus	Absorbed tetanus toxoid	1 to 3 doses	Born in Japan later than 1968 having immunity: once in ten years Born in Japan before 1967, or with no vaccination record: 3 doses for immunity, and once in every ten years (0, 3 to 8 weeks later, 12 to 18 months later)	JPY 6,100	10 years
Cholera	*Dukoral	2 to 3 doses	Children 2 to 5 years old: 3 doses by mouth at least one week (up to 6 weeks) apart, at least 3 weeks before the trip Age 6 years and older: 2 doses by mouth at least one week (up to 6 weeks) apart	JPY 12,000	2 years
Japanese encephalitis	Jebik V	3 doses	at 0, 1 to 4 weeks later, 1 year later	JPY 6,600	3 to 4 years
Meningococcal meningitis	Menactra	1 dose	only once (2 to 55 years old)	JPY 24,000	5 years
Yellow fever		1 dose	vaccinations are available only at quarantine stations	JPY 17,380	10 years

* Importation delivery cost (JPY 38,000 + tax) will be added to vaccine fee when it is imported.

Prophylaxis against	Product Name	Name of Drug	Dosing	Initiation	Discontinuation	Fee	Use in pregnancy
Malaria	Vibramycin tablet 100mg	Doxycycline	1 tablet orally, once daily	1 to 2 days (at the time before first exposure to malaria)	4 weeks (from the time after last exposure)	JPY 60/tablet	No; teratogenic
Malaria	Malarone tablet	Atovaquone (250mg) + Proguanil (100mg)	1 tablet orally, once daily	1 to 2 days (at the time before first exposure to malaria)	7 days (from the time after last exposure)	JPY 700/tablet	Insufficient data on use in pregnancy
Mountain sickness	Diamox tablet 250mg	Acetazolamide 250mg	0.5 tablet (one half tablet) twice daily, morning and evening	1 day before ascent	3 days after first dosing	JPY 100/tablet	Insufficient data on use in pregnancy

* Reservation and complete upfront payment including vaccine fee and initial consultation fee (JPY 2,880) is required.

※Tax included

* Vaccinations for children must be considered on together with their mandatory periodical vaccinations; individual consultation is required

* Regarding effective period of vaccine, refer JSTH (Japanese Society of Travel and Health)

* Regarding previously vaccinated rabies dosing interval, refer WHO position paper (April 2018)

January 2020